### REMARKS

Reconsideration and withdrawal of the rejections of the claims, in view of the amendments and remarks herein, is respectfully requested. Claims 1 and 50-53 are amended, and claims 56-57 are added; as a result, claims 1-31 and 44-57 are pending in this application.

Support for the amendments to claims 1 and 50-53, is found, for instance, at page 13, lines 19-25, page 19, lines 24-28, and page 20, lines 3-5 and 20-22 of the specification. Support for new claims 56 and 57 is found, for instance, in the example in the specification.

### The 35 U.S.C. § 112, First Paragraph, Rejections

The Examiner rejected claims 1, 8-9, 15-19, 23, 25, and 44-55 under 35 U.S.C. § 112, first paragraph, as lacking adequate written description. Specifically, the Examiner alleges that the specification teaches the structure of only a single representative species of SEQ ID Nos:2, 3 and 4, and fails to describe any other representative species by any identifying characteristics or properties other than the functionality of hybridizing to SEQ ID NO:2, 3 or 4 and to fails describe polypeptides [sic] having at least 80%, 85%, 90%, 95% or 97% sequence identity to SEQ ID NO:2, 3 or 4. The Examiner also asserts that there is no support for the primers or probe consisting of 15 to 40 nucleotides which include SEQ ID NO:2, 3 or 4. The Examiner also rejected claims 1, 8-9, 15-19, 23, 25, and 44-55 under 35 U.S.C. § 112, first paragraph, as containing new matter. In particular, the Examiner alleges that there is no support for probes or primers having about 15 to 40 nucleotides with at least 80% identity to SEQ ID NO:2, 3, or 4 or the complement thereof that hybridizes to SEQ ID NO:2, 3, or 4. These rejections are respectfully traversed.

The Examiner is respectfully reminded that <u>Applicant need not teach</u> what is well know to the art. vanA sequences, including vanA-specific probes and primers, <u>were well known to the</u> art, as were amplification and hybridization assays to detect those and other sequences (see Petrich et al., <u>Mol. Cell Probes</u>, <u>13</u>:275 (1999) and U.S. Patent No. 6,274,316; both of record, as well as the Table of Contents for <u>Diagnostic Molecular Microbiology</u>; <u>Principles and Applications</u>, Pershing, ed., American Society of Microbiology, 1993 (a copy is enclosed herewith).

Title: METHOD AND KIT FOR IDENTIFYING VANCOMYCIN-RESISTANT ENTEROCOCCUS

Moreover, Applicant's specification clearly identifies the nucleotide sequence corresponding to nucleotides 870 to 896, 851 to 868 and 898 to 917 of van (see Figure 1 and SEQ ID NOs. 2-4).

The Examiner cannot reasonably contend that the recitation of a primer or probe that forms a hybrid with, and has at least 80% nucleic acid sequence identity to, a particular nucleic acid structure does not convey a common structure or function. Hybrid formation and percent nucleic acid sequence identity between two nucleic acid molecules *clearly* convey a common structure (see, for example, the Table of Contents for <u>Diagnostic Molecular Microbiology</u>: <u>Principles and Applications</u>, Pershing, ed., American Society of Microbiology, 1993; <u>The Polymerase Chain Reaction</u>, Mullis et al., eds., Birkhauser, 1994 (copy enclosed), and <u>Molecular Cloning</u>: <u>A Laboratory Manual</u>, Sambrook et al., eds., Cold Spring Harbor Press, 2001 (copy enclosed)).

The Examiner also cannot reasonably contend that the skilled artisan cannot envision the detailed structure of nucleic acid sequences with at least 80% nucleic acid sequence identity to SEQ ID NO:2, 3 or 4. For instance, SEQ ID NO:2 has 18 nucleotides, SEQ ID NO:3 has 27 nucleotides, and SEQ ID NO:4 has 20 nucleotides. A sequence with 3 nucleotide substitutions relative to SEQ ID NO:2 has 83% identity thereto, a sequence with 6 nucleotide substitutions relative to SEQ ID NO:3 has 81% identity thereto, and a sequence with 4 nucleotide substitutions relative to SEQ ID NO:4 has 80% identity thereto

Moreover specifically, SEQ ID NO:2 corresponds to CCGGTGGCAGCTACGTTT (18

nucleotides). 80% of 18 is 14.4. Therefore, probes within the scope of invention include those with up to 3 nucleotide substitutions in SEQ ID NO:2. The following represents a set of variants of SEQ ID NO:2 with a single substitution (indicated by underlining):

CGGGTGGCAGCTACGTTT, CAGGTGGCAGCTACGTTT, and

CTGGTGGCAGCTACGTTT. The following represents a set of variants of SEQ ID NO:2 with two substitutions (indicated by underlining): CGGGTGGCAGCTAGGTTT, CAGGTGGCAG

CTAAGTTT, and CTGGTGGCAGCTATGTTT, and the following represents a set of variants of SEQ ID NO:2 with three substitutions (indicated by underlining): CGGGGGGCAG

CTAGGTTT. CAGGAGGCAGCTAAGTTT, and CTGGCGGCAGCTATGTTT.

Given that the Example in the specification discloses the use of SEQ ID Nos. 2-4 in a method to detect vanA sequences in a sample, and that the specification discloses that sequences

with close structural relatedness to SEQ ID Nos. 2-4, i.e., those with at least 80% nucleic acid sequence identity to the 18 nucleotides corresponding to SEQ ID NO;2, the 27 nucleotides corresponding to SEQ ID NO:3, and the 20 nucleotides corresponding to SEQ ID NO:4, are useful in the claimed methods, one of skill in the art would recognize that Applicant was in possession of the claimed invention.

It is unclear to Applicant why claims 54-55 do not satisfy the written description requirement, as they are directed to primers and probes with a specific sequence.

The fundamental inquiry with regard to new matter is whether the material added by amendment was inherently contained in the original application. *Litton Sys., Inc. v. Whirlpool Corp.*, 728 F.2d 1423, 1438, 221 U.S.P.Q. 97, 106 (Fed. Cir. 1984). The issue is not whether a specific new word of a claim was used in the specification as filed but whether the concept expressed by the word was present. *In re Anderson*, 471 F.2d 1237, 176 U.S.P.Q. 331 (C.C.P.A. 1973).

As filed, claim 1 recited that the *vanA*-specific oligonucleotide probe comprises sequences which include sequences substantially corresponding to nucleotides 870 to 896 of the *vanA* gene, the complement thereof, or a portion thereof, sequences substantially corresponding to nucleotides 851 to 868 of the *vanA* gene, the complement thereof, or a portion thereof, or sequences substantially corresponding to nucleotides 898 to 917 of the *vanA* gene, the complement thereof, or a portion thereof.

Claim 32 (as filed) is directed to an oligonucleotide composition comprising a first oligonucleotide comprising sequences substantially corresponding to nucleotides 870 to 896 of the *vanA* gene, the complement thereof, or a portion thereof, an oligonucleotide comprising sequences substantially corresponding to nucleotides 851 to 868 of the *vanA* gene the complement thereof, or a portion thereof, an oligonucleotide comprising sequences substantially corresponding to nucleotides 898 to 917 of the *vanA* gene, the complement thereof, or a portion thereof, or a combination thereof, wherein the oligonucleotide hybridizes under stringent hybridization conditions to *vanA* DNA. Claim 34 (as filed) depends on claim 32, and is directed to at least one oligonucleotide that has the length and sequence of any of SEO ID NOs:2-4.

Pages 10 and 11 of the specification disclose that:

An "oligonucleotide" is a polynucleotide having two or more nucleotide subunits covalently joined together.

A "primer" is a single-stranded polyoligonucleotide that combines with a complementary single-stranded target to form a double-stranded hybrid, which primer in the presence of a polymerase and appropriate reagents and conditions, results in nucleic acid synthesis.

A "probe" is a single-stranded polynucleotide that combines with a complementary single-stranded target polynucleotide to form a double-stranded hybrid.

Page 6 of the specification discloses:

In one embodiment, the oligonucleotides of the invention include sequences <u>substantially corresponding to nucleotides 851 to 868</u> of the <u>vanA</u> gene (SEQ ID NO.2; an exemplary <u>vanA</u> gene has SEQ ID NO:1 from <u>E. faecium pIP816 gi</u> 43335, also see Figure 1, Accession No. X56895 which corresponds to SEQ ID NO:11), or the complement thereof, or a portion thereof; sequences <u>substantially corresponding to nucleotides 870 to 896</u> of the <u>vanA</u> gene (SEQ ID NO:3), the complement thereof, or a portion thereof; sequences <u>substantially corresponding to nucleotides 890 to 917</u> of the <u>vanA</u> gene (SEQ ID NO:4), the complement thereof, or a portion thereof (emphasis added).

The Example discloses the use of primers having SEQ ID Nos. 2 and 4 to amplify vanAspecific sequences in a sample and a probe having SEQ ID NO:3 to detect vanA-specific sequences in a sample.

In the Amendment filed on May 30, 2007, Applicant indicated that support for the amendments to claim 1 is found, for instance, at page 13, lines 19-25, page 19, line 24-page 20, line 2, and page 20, lines 3-5 and 20-22 of the specification. Page 13 discloses that:

One skilled in the art will understand that probes or primers that substantially correspond to a reference sequence or region can vary from that reference sequence or region and still hybridize to the same target nucleic acid sequence. Probes of the present invention substantially correspond to a nucleic acid sequence or region if the percentage of identical bases or the percentage of perfectly complementary bases between the probe and its target sequence is from 100% to 80% or from 0 base mismatches in a 10 nucleotide target sequence to 2 bases mismatched in a 10 nucleotide target sequence. In one embodiment, the percentage is from 100% to 85%. In another embodiment this percentage is from 90% to 100%; and in yet other embodiments, this percentage is from 95% to 100% (emphasis added).

### Page 19 discloses that:

Preferred methods for detecting the presence of the vanA or vanB gene, include the step of contacting a test sample with at least two oligonucleotide primers under conditions that

preferentially amplify vanA and/or vanB sequences... While oligonucleotides probes of different lengths and base composition may be used for detecting the vanA gene or the vanB gene, preferred oligonucleotides have lengths from 15 up to 40 nucleotides and are sufficiently homologous to the target nucleic acid to permit amplification of a vanA or vanB template and/or hybridization to such a template under high stringency conditions (emphasis added).

### Page 19 also discloses that:

[T]he specific sequences described herein also may be provided in a nucleic acid cloning vector or transcript or other longer nucleic acid and still can be used for amplifying or detecting the vanA gene or the vanB gene, i.e., the probes may include sequences unrelated to the vanA or vanB gene, for instance at the 5' end, the 3' end, or both the 5' and 3' ends. Likewise, primers may include sequences unrelated to the vanA gene and/or the vanB gene, e.g., at the 5' end (emphasis added).

### Page 20 discloses that:

Preferred primers and probes have sequences of up to 40 nucleotides in length and preferably have at least 17 contiguous nucleotides corresponding to sequences in the vanA gene or the vanB gene, or the complement thereof... Preferably, the probes specifically hybridize to vanA or vanB DNA only under conditions of high stringency. Under these conditions only highly complementary nucleic acid hybrids will form (i.e., those having at least 14 out of 17 bases in a contiguous series of bases being complementary). Hybrids will not form in the absence of a sufficient degree of complementarity (emphasis added).

Thus, the specification clearly provides support for the recited probes and primers.

Accordingly, withdrawal of the § 112, first paragraph, rejections is respectfully requested.

### The 35 U.S.C. § 112, Second Paragraph, Rejections

The Examiner rejected claims 1, 8-9, 15-19, 23, 25, and 44-55 under 35 U.S.C. § 112, second paragraph, as being indefinite. In particular, the Examiner asserts that 1) claims 1 and 50-53 are unclear, as claims 1 and 50-53 recite the broad limitation of having 15 to 40 nucleotides, yet SEQ ID NOs:2, 3 and 4 do not have 40 nucleotides (the sequences have 18, 27 and 20 nucleotides, respectively), and the claims and specification fail to disclose what the other nucleotides are: 2) the term "hybridizes" in the claims is a relative term which renders the claims

Title: METHOD AND KIT FOR IDENTIFYING VANCOMYCIN-RESISTANT ENTEROCOCCUS

indefinite; and 3) the phrase "one which under the same conditions hybridizes" in claim 1 is a relative phrase which renders the claims indefinite. These rejections, as they may be maintained with respect to the pending claims, are respectfully traversed.

The claims no longer recite "one which under the same conditions hybridizes", thereby obviating the § 112(2) rejection thereof.

It is Applicant's position that the metes and bounds of the recited primers and probe are clear. That is, the probe is a vanA-specific oligonucleotide that forms a hybrid with vanA nucleic acid in a sample. The probe consists of 15 to 40 nucleotides and has a sequence with at least 80% nucleic acid sequence identity to SEO ID NO:3 or the complement of SEO ID NO:3. The probe is one which forms a hybrid with SEO ID NO:3 or its complement. Thus, the probe is of a particular length and has sequence homology to SEQ ID NO:3 or its complement.

The two primers are oligonucleotides that amplify vanA nucleic acid. The primers consist 15 to 40 nucleotides. One (the first) primer has a sequence with at least 80% nucleic acid sequence identity to SEQ ID NO:2, and the other (the second) primer has a sequence with at least 80% nucleic acid sequence identity to SEO ID NO:4. The first primer forms a hybrid with the complement of SEQ ID NO:2, and the second primer forms a hybrid with the complement of SEO ID NO:4. Hence, the primers are of a particular length and have sequence homology to SEQ ID NO:2 or SEQ ID NO:4.

For instance, a nucleic acid sequence which consists of 15 contiguous nucleotides of SEQ ID NO:2 (SEO ID NO:2 has 18 nucleotides) has a sequence with 15 nucleotides having 100% nucleic acid sequence identity to SEQ ID NO:2; a nucleic acid sequence which consists of 15 nucleotides, 14 of which are identical in sequence to those in SEQ ID NO:2 and one of which is different, has a sequence with 15 nucleotides having at least 80% nucleic acid sequence identity to SEQ ID NO:2; and a nucleic acid sequence which consists of 40 nucleotides, 15 of which are identical in sequences to those in SEQ ID NO:2, has a sequence with 15 nucleotides having 100% nucleic acid sequence identity to SEO ID NO:2.

Moreover, one of skill in the art understands that sequences other than those needed for specificity of a probe or primer may be included with the probe or primer, e.g., a restriction enzyme site may be included in a primer sequence to facilitate cloning of sequences amplified with the primer. For example at page 19 of the specification, it is disclosed that:

Title: METHOD AND KIT FOR IDENTIFYING VANCOMYCIN-RESISTANT ENTEROCOCCUS

the specific sequences described herein also may be provided in a nucleic acid cloning vector or transcript or other longer nucleic acid and still can be used for amplifying or detecting the vanA gene or the vanB gene, i.e., the probes may include sequences unrelated to the vanA or vanB gene, for instance at the 5' end, the 3' end, or both the 5' and 3' ends. Likewise, primers may include sequences unrelated to the vanA gene and/or the vanB gene, e.g., at the 5' end.

Therefore, the metes and bounds of the recited probe and primers is clear.

It is Applicant's position that one of skill in the art would understand the metes and bounds of the term "hybridizes" as it is conventionally used in the art. See, e.g., Petrich et al. supra, and U.S. Patent No. 6,274,316, supra, which disclose primers or probes that hybridize to vanA sequences in a biological sample. It is also Applicant's position that the selection of amplification and/or hybridization conditions specific for vanA sequences is conventional in the art. See, for instance, Petrich et al., supra, Patel et al., "Multiplex PCR Detection of vanA, vanB, vanC-1, and vanC-2/3 Genes in Enterococci", J. Clin. Microbiology, 35: 703 (1997); Petrich et al., "Effect of Routine Use of a Multiplex PCR for Detection of vanA- and vanB- Mediated Enterococcal Resistrance on Accuracy, Costs and Earlier Reporting", Diagnostic Microbiology and Infectious Disease, 41:215 (2001); and Satake et al., "Detection of Vancomycin-Resistant Enterococci in Fecal Samples by PCR", J. Clin. Microbiology, 35:2325 (1997); all of record.

Moreover, one of skill in the art is aware that more than one set of conditions can result in the specific hybridization of two nucleic acid sequences.

Even if, assuming for the sake of argument, the metes and bounds of the term "hybridizes" was not readily recognizable to one skilled in the art, the specification discloses particular conditions at page 20 (for amplification reactions with primers) and pages 24-25 (for probes). Thus, the scope of the claims would be clear to a person of skilled in the art, particularly when read in light of the specification.

Nevertheless, to advance the application, "hybridizes" has been deleted from claim 1.

Accordingly, withdrawal of the § 112, second paragraph, rejections is respectfully requested.

### CONCLUSION

Applicant respectfully submits that the claims are in condition for allowance, and notification to that effect is earnestly requested. The Examiner is invited to telephone Applicant's attorney at (612) 373-6959 to facilitate prosecution of this application.

If necessary, please charge any additional fees or credit overpayment to Deposit Account No. 19-0743.

Respectfully submitted,

SCHWEGMAN, LUNDBERG & WOESSNER, P.A. P.O. Box 2938

Minneapolis, MN £5402

(612) 373-6959

Date MW 5, 2008

Janet E. Hmbretson Reg. No. 39,665

CANDIS BUENDING

Name

Signature

## Diagnostic Molecular Microbiology

PRINCIPLES AND APPLICATIONS

Edited by

David H. Persing, M.D., Ph.D.

Section of Clinical Microbiology, Mayo Clinic, Rochester, Minnesota

Thomas F. Smith, Ph.D.

Section of Clinical Microbiology, Mayo Clinic, Rochester, Minnesota

Fred C. Tenover, Ph.D.

National Center for Infectious Diseases, Centers for Disease Control, Atlanta, Georgia

Thomas J. White, Ph.D.

Roche Molecular Systems, Alameda, California

AMERICAN SOCIETY FOR MICROBIOLOGY WASHINGTON, D.C.

Copyright © 1993 Mayo Foundation Rochester, MN 55905

Library of Congress Cataloging-in-Publication Data

Diagnostic molecular microbiology: principles and applications / edited by David H. Persing . . . [et al.].

p. cm. Includes index.

ISBN 1-55581-056-X

 DNA probes-Diagnostic use. 2. Diagnostic microbiology—Technique. 3. Nucleic acid hybridization. 4. Communicable diseases—Diagnosis—Laboratory manuals. 1. Persing, David H. QR69.04D5 1993

616.9'047583-dc20

92-38523 CIP All Rights Reserved
Printed in the United States of America

This book was coedited and chapter 1 was cowritten by Fred C. Tenover in his private capacity. No official support or endorsement by CDC is intended or should be inferred.

Cover illustration by Tomo Narashima

### **Contents**

|      | ibutors  | ix  |
|------|--|-----|
|      |  | xix |
|      | ord. Albert Balows   | xxi |
| refa | ce. David H. Persing   | XXI |
| art  | I. Principles of Diagnostic Molecular Microblology   |     |
| 1.   | Nucleic Acid Probes for Detection and Identification of Infectious Agents. Fred C. Tenover and Elizabeth R. Unger  | 3   |
| 2.   | Molecular Typing Methods. Bala Swaminathan and<br>Ghassan M. Matar   | 26  |
| 3.   | In Vitro Nucleic Acid Amplification Techniques.  David H. Persing  | 51  |
| 4.   | Target Selection and Optimization of Amplification Reactions.  David H. Persing  | 88  |
| 5.   | Amplification Product Inactivation Methods. David H. Persing and George D. Cimino  | 105 |
| 6.   | Sample Preparation Methods. Larry Greenfield and Thomas J. White   | 122 |
| 7.   | Amplification Product Detection Methods. Thomas J. White   | 138 |
| 8.   | Laboratory Design and Work Flow. Bruce J. McCreedy and<br>Thomas H. Callaway   | 149 |
|      | Quality Control of Polymerase Chain Reaction. Elizabeth A.   |     |
| 9.   | Dragon, Joanne P. Spadoro, and Roberta Madej   | 160 |
| art  | II. Applications of Diagnostic Molecular Microbiology  |     |
| ECT  | ION 1. Bacterial Pathogens   |     |
| 1.1  | Novel and Uncultured Pathogens. Philip Kirschner,  |     |
|      | Albrecht Meier, and Erik C. Böttger  | 173 |
| 1.2  | PCR Detection of Mycobacterium tuberculosis. Kathleen D. Eisenach, M. Donald Cave, and Jack T. Crawford  | 191 |
| 1.3  | PCR Detection of Rickettsia rickettsia and Ehrlichia chaffeensis.  Burt E. Anderson  | 197 |
| 1.4  | PCR Detection of Borrelia burgdorferi. Paul N. Rys   | 203 |
| 1.5  | Rapid Ribosequencing for the Detection and Classification of Tick-Borne Borrelia burgdorferi and Related Species.  Ulf B. Göbel, Barbara Graf, and Thomas Adam | 211 |
| 1.6  | PCR Detection of Yersinia pestis in Fleas. J. Hinnebusch and Tom G. Schwan   | 218 |
| 1.7  | PCR Detection of Treponema pallidum. Justin D. Radolf  | 224 |
| 1.8  | Molecular Detection of Treponema pallidum via a 16S Target.  | 230 |
|      | Jeffrey S. Sartin  |     |
| 1 9  | PCP Detection of Chlamydia trachomatis, Linda D. Bobo  | 433 |

Contents vi 1.10 3SR Detection of Chlamydia trachomatis. Paul V. Haydock 1.11 PCR Detection of Chlamydia pneumoniae. Lee Ann Campbell . . . . 247 1.12 PCR Detection of Mycoplasma pneumoniae. Maurice Garret 1.13 Rapid and Sensitive PCR Method for Identification of Mycoplasma Species in Tissue Culture, Jolene G. Wong-Lee 1.14 PCR Detection of Legionella pneumophila and L. dumoffii in 1.15 PCR Detection of Toxin Genes in Strains of Vibrio cholerae O1. Orjan Olsvik, Tanja Popovic, and Patricia I. Fields . . . . . . . . . . . . 1.16 PCR Detection of Heat-Stable, Heat-Labile, and Shiga-Like Toxin Genes in Escherichia coli, Ørian Olsvik and 1.17 PCR Detection of Shigella Species and Enteroinvasive 1.18 PCR Detection of Helicobacter pylori, Joan L. Valentine . . . . . . 282 1.19 PCR Detection of Toxigenic Clostridium difficile. Paul H. Gumerlock, Yajarayma J. Tang, and Joseph Silva, Jr. . . . . . . . 288 1.20 PCR Detection of Genes for Enterotoxins, Exfoliative Toxins, and Toxic Shock Syndrome Toxin-1 in Staphylococcus aureus. 1.21 PCR Detection of Bacteria Found in Cerebrospinal Fluid. SECTION 2. Viral Pathogens 2.1 PCR Detection of Human Immunodeficiency Virus Type 1 Proviral DNA Sequences. Shirley Kwok and John J. Sninsky. . . . . 2.2 PCR Detection of Human T-Cell Lymphotrophic Virus Type I. Moses Rodriguez, Naraporn Prayoonwiwat, and Larry R. Pease . . 316 2.3 PCR Detection of Hepatitis B Virus. Osamu Yokosuka, 2.4 Reverse Transcriptase-PCR for Hepatitis C Virus RNA. Judith C. Wilber, Pamela J. Johnson, and Mickey S. Urdea . . . . 327 2.5 PCR Detection of Herpes Simplex Virus DNA Sequences in Cerebrospinal Fluid. Mark J. Espy, Jaber Aslanzadeh, and 2.6 PCR Detection of Herpes Simplex Virus, Richard W. Cone and Ann C. Hobson.... PCR Detection and Typing of Epstein-Barr Virus. Amalio Telenti. 344 2.8 PCR Detection of Cytomegalovirus DNA Sequences in Clinical 2.9 PCR Detection of Varicella-Zoster Virus, 

| Content | is   | vii        |
|---------|--|------------|
| 2.12    | PCR Detection and Differentiation of Influenza Virus A, B, and C Strains. Wandong Zhang and David H. Evans   | 374        |
| 2.13    | PCR Detection of Rotavirus. Vera Gouvea  | 383        |
| 2.14    | PCR Detection of Human Adenoviruses. Michael McDonough,<br>Olen Kew, and John Hierholzer   | 389        |
| 2.15    | PCR Detection of Rubella Virus. Clark B. Inderlied and Kevin A. Nash   | 394        |
| 2.16    | PCR Detection of the Human Enteroviruses. José R. Romero and Harley A. Rotbart   | 401        |
| 2.17    | PCR Detection of Genital Human Papillomavirus. Heidi M. Bauer and M. Michele Manos.  | 407        |
| 2.18    | Rapid Amplification of Human Papillomavirus Type 16 and 18 E6 and E7 mRNA by 3SR. Janice T. Brown and Alan T. Wortman  | 414        |
| SECT    | FION 3. Fungal Pathogens   |            |
| 3.1     | A Model PCR/Probe System for the Identification of Fungal Pathogens. Barbara H. Bowman   | 423        |
| 3.2     | PCR Identification of Cryptococcus neoformans. T. G. Mitchell, E. Z. Freedman, W. Meyer, T. J. White, and J. W. Taylor   | 431        |
| 3.3     | PCR Detection of Pneumocystis carinii: Diagnosis and Therapeutic Monitoring. Yoshikazu Nakamura  | 437        |
| SEC1    | FION 4. Parasitic Pathogens  |            |
| 4.1     | PCR Detection of Trypanosoma cruzi, African Trypanosomes, and Leishmania Species. Louis V. Kirchhoff and John E. Donelson                                      | 443        |
| 4.2     | PCR Detection of Leishmania braziliensis. Jorge Arévalo, Rocio Inga, and Martin López  | 456        |
| 4.3     | PCR Detection of Plasmodium Species in Blood and Mosquitoes. Konstantinos Mathiopoulos, Madama Bouaré, Glenn McConkey, and Thomas McCutchan                    | 462        |
| 4.4     | PCR Detection of Pathogenic Entamoeba histolytica and<br>Differentiation from Other Intestinal Protozoa by   |            |
|         | Riboprinting. C. Graham Clark  | 468<br>475 |
| 4.5     | PCR Detection of Babesia microti. David H. Persing   | 480        |
| 4.6     | PCR Detection of Giardia lamblia. Judith B. Weiss  | 400        |
| SECT    | TION 5. Novel Organisms  |            |
| 5.1     | Universal Bacterial 16S rDNA Amplification and Sequencing.  David A. Relman  | 489        |
| 5.2     | PCR-Based Detection of the Uncultured Bacillus of Whipple's Disease. David A. Relman   | 496        |
| 5.3     | Detection of Human Herpesvirus 6 and Human Herpesvirus 7 by PCR Amplification. Mary E. Klotman, Paolo Lusso, David Bacchus, Mario Corbellino, Ruth F. Jarrett, |            |
|         | and Zwi N. Berneman  | 501        |

| viii       | `   | Contents |
|------------|---|----------|
| 5.4        | PCR Detection of Mycoplasma fermentans Infection in Blood and Urine, Richard YH. Wang and Shyh-Ching Lo   | . 511    |
| SEC        | TION 6. Antimicrobial Resistance Loci   |          |
| 6.1        | Detection of Rifampin Resistance Mutations in Mycobacterium tuberculosis and M. leprae. Paul Imboden, Stewart Cole,   | 610      |
| 6.2        | Thomas Bodmer, and Amalio Telenti PCR Detection of Human Immunodeficiency Virus Drug Resistance Mutations. Brendan A. Larder and Charles A. B. Boucher                                |          |
| 6.3        | PCR Detection of erm Erythromycin Resistance Genes by<br>Using Degenerate Oligonucleotide Primers. Michel Arthur,<br>Catherine Molinas, and Claude Mabilat                            |          |
| 6.4        |   |          |
| 6.5        | PCR Detection of Penicillinase-Producing Neisseria gonorrhoeae.  Jean-Luc Simard and Paul H. Roy  | . 543    |
| 6.6        | PCR Detection of Genes Coding for Aminoglycoside-Modifying Enzymes. Jos A. M. van de Klundert and John S. Vliegenthart  | . 547    |
| 6.7        | PCR Detection and Identification of Genes for Extended-Spectrum<br>8-Lactamases. Claude Mabilat and Sylvie Goussard   | . 553    |
| SEC        | FION 7. Molecular Typing Methods  |          |
| 7.1        | Application of Pulsed-Field Gel Electrophoresis to Molecular Epidemiology. Joel N. Maslow, Alexander M. Slutsky,  |          |
| 7.2        | and Robert D. Arbeit  | . 563    |
|            | Cheryl A. Bopp, Orjan Olsvik, and Julia A. Kiehlbauch   |          |
| 7.3        | Chemiluminescent Ribotyping. Cynthia A. Gustaferro  |          |
| 7.4<br>7.5 | PCR Analysis of Integrons, Céline Lévesque and Paul H. Roy<br>Characterization of Pathogenic Microorganisms by Genomic<br>Fingerprinting Using Arbitrarily Primed PCR. John Welsh and | . 590    |
|            | Michael McClelland  | . 595    |
| SEÇT       | FION 8. Methods   |          |
| 8.1        | Simple and Rapid Sample Preparation Methods for Whole Blood and Blood Plasma. Lily Lin, Yu Gong, Ken Metchette, George D. Chnino, John E. Hearst, and Stephen T. Isaacs               | . 605    |
| 8.2        | Universal Bacterial DNA Isolation Procedure. L. M. Graves and B. Swaminathan  |          |
| Anner      | dix. Suggested Materials. Paul N. Rys   | 623      |
|            | unit. Suggested Materials, Full 14, Nys   |          |

# The Polymerase Chain Reaction

Kary B. Mullis François Ferré Richard A. Gibbs Editors

Foreword by James D. Watson

With 112 Illustrations

Kary B. Mullis La Jolla, CA 92037 USA

François Ferré
The Immune Response Corporation
5935 Darwin Court
Carlsbad, CA 92008
USA

Richard A. Gibbs Institute for Molecular Genetics Baylor College of Medicine One Baylor Plaza Houston, TX 77030 USA

Printed on acid-free paper.

© 1994 Birkhäuser Boston

Birkhäuser

Copyright is not claimed for works of U.S. Government employees.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the copyright owner.

The use of general descriptive names, trademarks, etc., in this publication, even if the former are not especially identified, is not to be taken as a sign that such names, as understood by the Trade Marks and Merchandise Marks Act, may accordingly be used freely by anyone.

While the advice and information in this book are believed to be true and accurate at the date of going to press, neither the authors nor the editors nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Permission to photocopy for internal or personal use, or the internal or personal use of specific clients, is granted by Birkhäuser Boston for libraries and other users registered with the Copyright Clearance Center (CCC), provided that the base fee of \$6.0 per copy, pais \$0.20 per page is paid directly to CCC, 21 Congress Street, Salem, MA 01970, USA. Special requests should be addressed directly to Birkhäuser Boston, 675 Masachusetts Avenue, Cambridge, MA 02193, USA.

ISBN 0-8176-3607-2 (hardcover) ISBN 0-8176-3750-8 (softcover) ISBN 3-7643-3750-8 (softcover) ISBN 3-7643-3750-8 (softcover)

Typeset by Sherman Typography, York, PA.
Printed and bound by The Maple Press Company, York, PA.
Printed in the United States of America.

987654321

Cover illustration: Based on a photograph taken by Kary Mullis in a museum in Cologne. "The mostic had been taken from Italy by the Germans. I think it was from Pompeii. I saw a similar design still lineat in its original location in Herculaneum but had no eamers that day. Veasuvius covered both clieta in 79 A.D. I would guess that the structure of DNA was probably worked out about two flousantly search golden was born and therefore nineten hundred and elighty-eight years before Crick. The Romans, however, seemed to think that atoms were square, and most likely the significance of the structure did escape their notice. Of course, this was long before Avery."

### Contents

| Foreword           |  |     |
|--------------------|--|-----|
| James D.           | . Watson   | v   |
| Preface<br>Kary B. | Mullis   | ix  |
| Contribu           | tors   | xix |
|                    | PART ONE: METHODOLOGY  |     |
| I. Basi            | c Methodology  |     |
|                    | Manipulation of DNA by PCR Kenshi Hayashi  | 3   |
|                    | Cloning PCR Products Michael A. Frohman.   | 14  |
|                    | Optimization of Multiplex PCRs  Jeffrey S. Chamberlain and Joel R. Chamberlain   | 38  |
|                    | Preparation of Nucleic Acids for Archival Material  Darryl Shibata   | 47  |
|                    | PCR Amplification of Viral DNA and Viral Host Cell mRNAs in Situ<br>Janet Embretson, Katherine Staskus, Ernest Retzel, Ashley T. Haase,<br>and Peter Bitterman | 55  |
| II. Qua            | ntitation  |     |
|                    | Quantitative PCR: An Overview F. Ferré, A. Marchese, P. Pezzoli, S. Griffin, E. Buxton, and V. Boyer   | 67  |
| 7.                 | Quantification of DNAs by the Polymerase Chain Reaction Using an<br>Internal Control   |     |
|                    | Donald M. Com  |     |

|     | 8    | i. RT-PCR and mRNA Quantitation Jamel Chelly and Axel Kahn  | . 97 |
|-----|------|---|------|
|     | 9    | . Analysis of Human T-Cell Repertoires by PCR  Dominic G. Spinella and James M. Robertson   |      |
| Ш   | . No | onisotopic Detection  |      |
|     | 10   | . Ultrasensitive Nonradioactive Detection of PCR Reactions: An Overview Richard H. Tullis   | 123  |
|     | 11   | Fluorescent Detection Methods for PCR Analysis Michael J. Heller  |      |
|     | 12   | Enzyme-Labeled Oligonucleotides  Eugene Tu and Edward Jablonski   | 142  |
|     | 13   | Application of the Hybridization Protection Assay (HPA) to PCR<br>Norman C. Nelson and Sherrol H. McDonough   |      |
| IV. | Ins  | trumentation  |      |
|     | 14.  | PCR Instrumentation: Where Do We Stand? Christian C. Oste   | 165  |
|     | 15.  | Rapid Cycle DNA Amplification Carl T. Wittwer, Gudrun B. Reed, and Kirk M. Ririe  | 174  |
|     | 16.  | Automating the PCR Process H.R. Garner  | 182  |
| v.  | Sec  | quencing  |      |
|     | 17.  | PCR and DNA Sequencing Bjorn Andersson and Richard A. Gibbs.  | 201  |
|     | 18.  | Phage Promoter-Based Methods for Sequencing and Screening for<br>Mutations  |      |
|     | 19   | Steve S. Sommer and Erica L. Vielhaber  Capture PCR: An Efficient Method for Walking Along Chromosomal  | 214  |
|     |      | DNA and cDNA  |      |
|     |      | André Rosenthal, Matthias Platzer, and D. Stephen Charnock-Jones  | 222  |
|     |      | PART TWO: APPLICATIONS  |      |
| I.  | Gen  | eral Applications   |      |
|     | 20.  | In Vitro Evolution of Functional Nucleic Acids: High-Affinity RNA<br>Ligands of the HIV-1 rev Protein<br>Craig Tuerk, Sheela MacDougal-Waugh, Gerald Z. Hertz, and Larry Gold |      |
|     | 21.  | The Application of PCR to Forensic Science Bruce Budowle, Antii Sajantila, Manfred N. Hochmeister, and Catherine T. Comey   | 233  |
|     | 22.  | Recreating the Past by PCR Matthias Höss, Oliva Handt, and Svante Pääbo   | 257  |
|     | 23.  | Nonbiological Applications<br>Gavin Dollinger   | 265  |

| п.    | Ge  | Genetic Analysis  |    |  |  |  |
|-------|-----|---|----|--|--|--|
|       | 24. | RI-PCR and Gene Expression Didier Montarras, Christian Pinset, Jamel Chelly, and Axel Kahn  | 27 |  |  |  |
|       | 25. | Fingerprinting Using Arbitrarily Primed PCR: Application to Genetic Mapping, Population Biology, Epidemiology, and Detection of Differentially Expressed RNAs  John Welsh and Michael McClelland. | 29 |  |  |  |
|       | 26. | Genetics, Plants, and the Polymerase Chain Reaction Bruno W.S. Sobral and Rhonda J. Honeycutt   | 30 |  |  |  |
| III.  | Ass | sessment of Therapy Effectiveness   |    |  |  |  |
|       | 27. | PCR Assessment of the Efficacy of Therapy in Philadelphia<br>Chromosome-Positive Leukemias<br>Stephen P. Hunger and Michael L. Cleary   | 32 |  |  |  |
|       | 28. | The Detection of Minimal Residual Disease (MRD) in Acute Lymphoblastic Leukemia Using Clone-Specific Probes Directed against V(D)J Junctional Sequences   |    |  |  |  |
|       |     | Luc d'Auriol and François Sigaux  | 33 |  |  |  |
|       | 29. | Assessment of Therapy Effectiveness: Infectious Disease Salvatore J. Arrigo   | 34 |  |  |  |
|       | 30. | Gene Therapy Richard A. Morgan and W. French Anderson   | 35 |  |  |  |
| IV.   | Dia | gnostics  |    |  |  |  |
|       | 31. | PCR and Cancer Diagnostics: Detection and Characterization of Single<br>Point Mutations in Oncogenes and Antioncogenes<br>Manuel Perucho  | 36 |  |  |  |
|       | 32. | Clinical Applications of the Polymerase Chain Reaction  |    |  |  |  |
|       |     | Belinda J.F. Rossiter and C. Thomas Caskey  | 39 |  |  |  |
|       | 33. | Infectious Diseases W. John Martin  | 40 |  |  |  |
|       |     | PART THREE: PCR AND THE WORLD OF BUSINESS   |    |  |  |  |
|       | 34. | PCR in the Marketplace Ellen Daniell  | 42 |  |  |  |
|       | 35. | PCR and Scientific Invention: The Trial of DuPont vs. Cetus Kary B. Mullis  | 42 |  |  |  |
|       |     | , p.  | 42 |  |  |  |
| Y 4 . |     |   |    |  |  |  |

VOLUME 1

# Molecular Cloning

A LABORATORY MANUAL

### THIRD EDITION

www.MolecularCloning.com

### Joseph Sambrook

PETER MACCALLUM CANCER INSTITUTE AND THE UNIVERSITY OF MELBOURNE, AUSTRALIA

### David W. Russell

UNIVERSITY OF TEXAS SOUTHWESTERN MEDICAL CENTER, DALLAS



# Molecular Cloning

A LABORATORY MANUAL

THIRD EDITION

www.MolecularCloning.com

### Joseph Sambrook

PETER MACCALLUM CANCER INSTITUTE AND THE UNIVERSITY OF MELBOURNE, AUSTRALIA

### David W. Russell

UNIVERSITY OF TEXAS SOUTHWESTERN MEDICAL CENTER, DALLAS



COLD SPRING HARBOR LABORATORY PRESS Cold Spring Harbor, New York

VOLUME 3

# Molecular Cloning

A LABORATORY MANUAL

### THIRD EDITION

www.MolecularCloning.com

### Joseph Sambrook

PETER MACCALLUM CANCER INSTITUTE AND THE UNIVERSITY OF MELBOURNE, AUSTRALIA

### David W. Russell

UNIVERSITY OF TEXAS SOUTHWESTERN MEDICAL CENTER, DALLAS



### Molecular Cloning

THIRD EDITION

©2001 by Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York All rights reserved

Printed in the United States of America

Front cover (paperback): The gene encoding green fluorescent protein was cloned from Aequorea victoria, a jellyfish found in abundance in Puget Sound, Washington State. This picture of a 50-mm medusa was taken on color film by flash photography and shows light reflected from various morphological features of the animal. The small bright roundish blobs in the photograph are symbiotic amphibods living on or in the medusa. The bright ragged area in the center is the jellyfish's mouth.

Bioluminescence from Aequarea is emitted only from the margins of the meduase and cannot be seen in this image. Bioluminescence of Aequarea, as in most species of jellyfish, does not look like a soft overall glow, but occurs only at the tim of the bell and, glowin the right viewing conditions, would appear as a string of nearly microscopic fusiform green lights. The primary luminescence produced by Aequarea is actually builth in color and is emitted by the protein aequorin. In a living jellyfish, light is emitted via the coupled green fluorescent protein, which causes the tuminescence to protein, which causes the tuminescence to protein, but a protein green green to the observer.

The figure and legend were kindly provided by Claudia Mills of the University of Washington, Friday Harbor. For further information, please see Mills, C.E. 1999-2000. Bioluminescence of Acquerea, a hydromedua. Electronic Internet document available at http://faculty.washington.edu/com/lils/Acquerea/thml. Published by the author, web page established June 1999, last updated 23 August 2009.

Back cover (paperback): A portion of a human cDNA array hybridized with a red fluor-tagged experimental sample and a green fluortagged reference sample. Please see Appendix 10 for details. (Image provided by Vivek Mittal and Michael Wigler, Cold Spring Harbor Laboratory.)

Library of Congress Cataloging-in-Publication Data

Sambrook, Joseph.

Molecular cloning: a laboratory manual / Joseph Sambrook, David W.

Russell .-- 3rd ed.

p. ; cm.

Includes bibliographical references and index.

ISBN 0-87969-576-5 (cloth) -- ISBN 0-87969-577-3 (pbk)

1. Molecular cloning--Laboratory manuals.

[DNLM: 1. Cloning, Molecular--Laboratory Manuals. QH 440.5 S187m

2001] I. Russell, David W. (David William), 1954- . II. Title.

QH442.2 .\$26 2001

572.8-\*dc21

00-064380

#### 10 9 8 7 6 5 4 3 2 1

People using the procedures in this manual do so at their own risk. Cold Spring Harbor Laboratory makes no representations or warranties with respect to the material set forth in this manual and has no liability in connection with the use of these materials.

All World Wide Web addresses are accurate to the best of our knowledge at the time of printing.

Certain experimental procedures in this manual may be the subject of national or local legislation or agency restrictions. Users of this manual are responsible for obtaining the relevant permissions, certificates, or licenses in these cases. Neither the authors of this manual nor Cold Spring Harbor Laboratory assume any reasonability for failure of a user to do so.

The polymerase child reaction process and other techniques in this manual may be or are covered by certain patent and proprietary right. User of this manual are reproposited for obtaining any ideosa necessary to practice PCR and other techniques or to commenciable the results of oth use C.O.D. SPRING HAR-BOR LASORATORY MAKES NO REPRESENTATION THAT USE OF THE INFORMATION IN THIS MANUAL WILL NOT INFRINGE ANY PATENT OR OTHER PROPRIETARY RIGHT.

Authorization to photocopy items for internal or personal sac, or the internal or personal use of specific clients, is granted by Cold Spring inhabitor laboratory. Press, provided that the appropriate Ees is paid directly to the Copyright Clearance Carner (CCC), Whise or all CCC at 222 Stotewood Dive, Dameers, MA 01923 (564-750-6400) for information about fees and regulations. Pitor to photocopying items for educational disastroom use, contact CCC at the aborted address. Additional information on CCC can be obtained at CCC Online as https://www.copyright.com.

All Cold Spring Harbor Laboratory Press publications may be ordered discoly from Cold Spring Harbor Laboratory Press, 10 Skyline Drive, Plainview, New York 1180:1290. Phone: 1-800-853-4388 in Continental U.S. and Camada. All other locations; (316) 340-1930. FNX: (316) 349-1946. Final Approximation for a complex catalog of all Cold Spring farbor Laboratory Press publications, with our World Work Shi Schizglich-Worked-Blory!

For orders from Purope, the Middle East, and Africa. British pound pricing is provided. Orders are foligiled and shipped from Cold Spring Harboy Laborauxy Press-Burope cite Lavis Marketing, 73 Line Walk, Headington, Oxford OXS 7AD, U.K. Phone: +44 (0) 1865 741541. FAX: +044 (0)1865 750079. E-mail: chippers.curope@ballo.jog. World Work Web Silte https://www.chipters.cu.pu/eb.

## Contents

| Preface           | xxi   |
|-------------------|-------|
| Acknowledgments   | xxiii |
| On-line Edition   | XXV   |
| Quotation Credits | vovii |

### Volume 1

| Chapter 1   |        |  |     |
|-------------|--------|--|-----|
| Plasmids an | d Thei | r Usefulness in Molecular Cloning  | 1.  |
|             | INITOC | DDUCTION   |     |
|             |        |  |     |
|             | PROT   | ocors  |     |
|             | In     | ntroduction to Preparation of Plasmid DNA by Alkaline Lysis with SDS (Protocols 1-3)   | 1.3 |
|             | 1      | Preparation of Plasmid DNA by Alkaline Lysis with SDS: Minipreparation   | 1.3 |
|             | 2      | Preparation of Plasmid DNA by Alkaline Lysis with SDS: Midipreparation   | 1.3 |
|             | 3      | Preparation of Plasmid DNA by Alkaline Lysis with SDS: Maxipreparation   | 1.3 |
|             | In     | stroduction to Preparation of Plasmid DNA by Boiling Lysis (Protocols 4 and 5)   | 1.4 |
|             | 4      | Preparation of Plasmid DNA by Small-scale Boiling Lysis  | 1.4 |
|             | 5      | Preparation of Plasmid DNA by Large-scale Boiling Lysis  | 1.4 |
|             | 6      | Preparation of Plasmid DNA: Toothpick Minipreparation  | 1.5 |
|             | 7      | Preparation of Plasmid DNA by Lysis with SDS   | 1.5 |
| *           | 8      | Purification of Plasmid DNA by Precipitation with Polyethylene Glycol  | 1.5 |
|             | 9      | Purification of Plasmid DNA by Chromatography  | 1.6 |
|             | 10     | Purification of Closed Circular DNA by Equilibrium Centrifugation in CsCl-Ethidium   | 1.6 |
| *           |        | Bromide Gradients: Continuous Gradients  |     |
|             | 11     | Purification of Closed Circular DNA by Equilibrium Centrifugation in CsCl-Ethidium<br>Bromide Gradients: Discontinuous Gradients | 1.6 |
|             | 12     | Removal of Ethidium Bromide from DNA by Extraction with Organic Solvents   | 1.7 |
|             | 13     | Removal of Ethidium Bromide from DNA by Ion-exchange Chromatography  | 1.7 |
|             | 14     | Removal of Small Fragments of Nucleic Acid from Preparations of Plasmid DNA by   | 1.7 |
|             |        | Centrifugation through NaCl  |     |
|             | 15     | Removal of Small Fragments of Nucleic Acid from Preparations of Plasmid DNA by   | 1.8 |
|             |        | Chromatography through Sephacryl S-1000  |     |
|             | 16     | Removal of Small Fragments of Nucleic Acid from Preparations of Plasmid DNA by<br>Precipitation with Lithium Chloride            | 1.8 |
|             | 17     | Directional Cloning into Plasmid Vectors   | 1.8 |

### Contents

| 18                 | Attaching Adaptors to Protruding Termini   | 1.88           |
|--------------------|--|----------------|
| 19                 | Blunt-ended Cloning into Plasmid Vectors   | 1,90           |
| 20                 | Dephosphorylation of Plasmid DNA   | 1.93           |
| 21                 | Addition of Synthetic Linkers to Blunt-ended DNA   | 1.98           |
| 22                 | Ligating Plasmid and Target DNAs in Low-melting-temperature Agarose  | 1.103          |
| 23                 | The Hanahan Method for Preparation and Transformation of Competent E. coli:<br>High-efficiency Transformation  | 1.105          |
| 24                 |  | 1.112          |
| 25                 | Preparation and Transformation of Competent E. coli using Calcium Chloride   | 1.116          |
| 26                 | Transformation of E. coli by Electroporation   | 1.119          |
| 27                 | Screening Bacterial Colonies Using X-gal and IPTG: α-Complementation   | 1.123          |
|                    | Alternative Protocol: Direct Application of X-gal and IPTG to Agar Plates  | 1.125          |
| 28                 | Screening Bacterial Colonies by Hybridization: Small Numbers   | 1.126          |
| 29                 | Screening Bacterial Colonies by Hybridization: Intermediate Numbers  | 1.129          |
|                    | Alternative Protocol: Rapid Lysis of Colonies and Binding of DNA to Nylon Filters  | 1,131          |
| 30                 | The second secon | 1.132          |
| 31                 |  | 1.135          |
| 32                 |  | 1.138          |
| INFOR              | MATION PANELS  |                |
| Chl                | oramphenicol   | 1.143          |
|                    | amycins  | 1.145          |
|                    | 3322   | 1.146          |
|                    | acycline   | 1.147          |
|                    | picillin and Carbenicillin   | 1.148<br>1.149 |
| X-ga               | omplementation   | 1,149          |
|                    | dium Bromide   | 1.150          |
|                    | idensing and Crowding Reagents   | 1.152          |
|                    | fication of Plasmid DNA by PEG Precipitation   | 1.152          |
|                    | ozymes   | 1.153          |
|                    | vethylene Glycol   | 1.154          |
| Ces                | ium Chloride and Cesium Chloride Equilibrium Density Gradients   | 1.154          |
| DN                 | A Ligases  | 1.157          |
|                    | ptors  | 1.160          |
| Elec               | rtroporation   | 1.162          |
| Chapter 2          |  |                |
| Bacteriophage λ ar | nd Its Vectors   | 2.1            |
|                    |  |                |
|                    | DUCTION  |                |
| PROTO              | OCOL\$   |                |
| . 1                | Plating Bacteriophage \( \lambda \)  | 2.25           |
|                    | • Additional Protocol: Plaque-Assay of Bacteriophages That Express β-Galactosidase   | 2.30           |
|                    | Additional Protocol: Macroplaques  | 2.31           |
|                    |  |                |
|                    |  |                |

|       | Conte  | nts vii |
|-------|--|---------|
|       |  | 2.32    |
| 2     |  | 2.34    |
| 3     | , , ,  | 2.34    |
|       | <ul> <li>Alternative Protocol: Preparing Stocks of Bacteriophage λ by Plate Lysis and<br/>Scraping</li> </ul>  |         |
| 4     | Preparing Stocks of Bacteriophage λ by Small-scale Liquid Culture  | 2.38    |
| 5     | Large-scale Growth of Bacteriophage λ: Infection at Low Multiplicity   | 2.40    |
|       | <ul> <li>Alternative Protocol: Large-scale Growth of Bacteriophage λ: Infection at High<br/>Multiplicity</li> </ul>                                      | 2.42    |
|       | Precipitation of Bacteriophage λ Particles from Large-scale Lysates  | 2.43    |
| 7     | Assaying the DNA Content of Bacteriophage & Stocks and Lysates by Gel<br>Electrophoresis   | 2.45    |
| 8     | Purification of Bacteriophage $\lambda$ Particles by Isopycnic Centrifugation through CsCl Gradients   | 2.47    |
|       | <ul> <li>Alternative Protocol: Purification of Bacteriophage λ Particles by Isopycnic<br/>Centrifugation through CsCl Equilibration Gradients</li> </ul> | 2.51    |
| 9     | Purification of Bacteriophage & Particles by Centrifugation through a Glycerol Step<br>Gradient  | 2.52    |
| 10    | Purification of Bacteriophage λ Particles by Pelleting/Centrifugation  | 2.54    |
| 11    |  | 2.56    |
| 12    |  | 2.59    |
| 13    | • • •  | 2.61    |
|       | Use as a Cloning Vector  |         |
| 14    | Preparation of Bacteriophage & DNA Cleaved with Two Restriction Enzymes for Us   | e 2.64  |
|       | as a Cloning Vector  |         |
|       | Alkaline Phosphatase Treatment of Bacteriophage λ Vector DNA   | 2.68    |
|       | Purification of Bacteriophage \( \lambda \) Arms: Centrifugation through Sucrose Density<br>Gradients  | 2.71    |
|       | Partial Digestion of Eukaryotic DNA for Use in Genomic Libraries: Pilot Reactions  | 2.76    |
| 18    | Partial Digestion of Eukaryotic DNA for Usé in Genomic Libraries: Preparative<br>Reactions   | 2.80    |
| 19    | Ligation of Bacteriophage λ Arms to Fragments of Foreign Genomic DNA   | 2.84    |
| 20    | Amplification of Genomic Libraries   | 2.87    |
| 21    | Transfer of Bacteriophage DNA from Plaques to Filters  | 2.90    |
|       | <ul> <li>Alternative Protocol: Rapid Transfer of Plaques to Filters</li> </ul>   | 2.95    |
| 22    | Hybridization of Bacteriophage DNA on Filters  | 2.96    |
| 23    | Rapid Analysis of Bacteriophage \( \) Isolates; Purification of \( \) DNA from<br>Plate Lysates  | 2.101   |
|       | Additional Protocol: Removing Polysaccharides by Precipitation with CTAB   | 2.105   |
| 24    | Rapid Analysis of Bacteriophage $\lambda$ Isolates: Purification of $\lambda$ DNA from Liquid Cultures   | 2.106   |
| INFOR | MATION PANELS  |         |
| Bac   | teriophages: Historical Perspective  | 2.109   |
|       | imizing Damage to Large DNA Molecules  | 2.110   |
| ln V  | fitro Packaging  | 2.110   |

viii Contents

| viii | Contents             |  |      |
|------|----------------------|--|------|
| Cha  | pter 3               |  |      |
| Wo   | rking with           | Bacteriophage M13 Vectors  | 3.1  |
|      |                      | INTRODUCTION   |      |
|      |                      | PROTOCOLS  | 3.17 |
|      |                      | 1 Plating Bacteriophage M13  | 3.20 |
|      |                      | <ol> <li>Growing Bacteriophage M13 in Liquid Culture</li> <li>Preparation of Double-stranded (Replicative Form) Bacteriophage M13 DNA</li> </ol> | 3.23 |
|      |                      | 4 Preparation of Single-stranded Bacteriophage M13 ONA   | 3.26 |
|      |                      | 5 Large-scale Preparation of Single-stranded and Double-stranded Bacteriophage   | 3.30 |
|      |                      | M13 DNA  |      |
|      |                      | 6 Cloning into Bacteriophage M13 Vectors   | 3.33 |
|      |                      | 7 Analysis of Recombinant Bacteriophage M13 Clones   | 3.39 |
|      |                      | <ul> <li>Alternative Protocol: Screening Bacteriophage M13 Plaques by Hybridization</li> </ul>   | 3.41 |
|      |                      | 8 Producing Single-stranded DNA with Phagemid Vectors  | 3.42 |
|      |                      | INFORMATION PANELS   |      |
|      |                      | Growth Times   | 3.49 |
|      |                      | Polyethylene Glycal  | 3.49 |
| a.   |                      |  |      |
|      | pter 4<br>rking with | High-capacity Vectors  | 4.1  |
|      |                      | INTRODUCTION   |      |
|      |                      |  |      |
|      |                      | PROTOCOLS  |      |
|      |                      | 1 Construction of Genomic DNA Libraries in Cosmid Vectors  | 4.11 |
|      |                      | 2 Screening an Unamplified Cosmid Library by Hybridization: Plating the Library onto Filters   | 4.24 |
|      |                      | <ul> <li>Additional Protocol: Reducing Cross-hybridization</li> </ul>  | 4.27 |
|      |                      | 3 Amplification and Storage of a Cosmid Library: Amplification in Liquid Culture   | 4.28 |
|      |                      | 4 Amplification and Storage of a Cosmid Library: Amplification on Filters  | 4.31 |
|      |                      | Alternative Protocol: Amplification on Plates  | 4.34 |
|      |                      | 5 Working with Bacteriophage P1 and Its Cloning Systems  | 4.35 |
|      |                      | Additional Protocol: Purification of High-molecular-weight DNA by Drop Analysis  | 4.44 |
|      |                      | <ul> <li>Alternative Protocol: Purification of High-molecular-weight Circular DNA by<br/>Chromatography on Qiagen Resin</li> </ul>               | 4.45 |
|      |                      | 6 Transferring P1 Clones between E, coli Hosts   | 4.46 |
|      |                      | 7 Working with Bacterial Artificial Chromosomes  | 4.48 |
|      |                      | 8 Isolation of BAC DNA from Small-scale Cultures   | 4.53 |
|      |                      | 9 Isolation of BAC DNA from Large-scale Cultures   | 4.55 |
|      |                      | 10 Working with Yeast Artificial Chromosomes   | 4.58 |
|      |                      | 11 Growth of S. cerevisiae and Preparation of DNA  | 4.67 |
|      |                      | 12 Small-scale Preparations of Yeast DNA   | 4.70 |
|      |                      | 13 Analyzing Yeast Colonies by PCR   | 4.72 |
|      |                      |  |      |

|                              | Coni   | tents ix |
|------------------------------|--|----------|
|                              | 14 Isolating the Ends of Genomic DNA Fragments Cloned in High-capacity Vectors:<br>Vectorette Polymerase Chain Reactions | 4.74     |
|                              | INFORMATION PANELS   |          |
|                              | Cre-loxP   | 4.82     |
|                              | Large-fragment Cloning Products and Services   | 4.86     |
| Chapter 5                    |  |          |
| Gel Electrop<br>Gel Electrop | horesis of DNA and Pulsed-field Agarose<br>horesis   | 5.1      |
|                              |  |          |
|                              | INTRODUCTION   |          |
|                              | PROTOCOLS  |          |
|                              | 1 Agarose Gel Electrophoresis  | 5.4      |
|                              | 2 Detection of DNA in Agarose Gels   | 5.14     |
|                              | 3 Recovery of DNA from Agarose Gels: Electrophoresis onto DEAE-cellulose Membranes                                       | 5.18     |
|                              | Recovery of DNA from Agarose and Polyacrylamide Gels: Electroelution into Dialysis Bags                                  | 5.23     |
|                              | 5 Purification of DNA Recovered from Agarose and Polyacrylamide Gels by<br>Anion-exchange Chromatography                 | 5.26     |
|                              | 6 Recovery of DNA from Low-melting-temperature Agarose Gels: Organic Extraction  | n 5.29   |
|                              | <ul> <li>Alternative Protocol: Recovery of DNA from Agarose Gels Using Glass Beads</li> </ul>                            | 5.32     |
|                              | 7 Recovery of DNA from Low-melting-temperature Agarose Gels: Enzymatic<br>Digestion with Agarase                         | 5.33     |
|                              | 8 Alkaline Agarose Gel Electrophoresis   | 5.36     |
|                              | <ul> <li>Additional Protocol: Autoradiography of Alkaline Agarose Gels</li> </ul>  | 5.39     |
|                              | 9 Neutral Polyacrylamide Gel Electrophoresis   | 5.40     |
|                              | 10 Detection of DNA in Polyacrylamide Gels by Staining   | 5.47     |
|                              | 11 Detection of DNA in Polyacrylamide Gels by Autoradiography  | 5.49     |
| \                            | 12 Isolation of DNA Fragments from Polyacrylamide Gels by the Crush and Soak Method                                      | 5.51     |
|                              | Introduction to Pulsed-field Gel Electrophoresis (Protocols 13-20)   | 5.55     |
| `                            | 13 Preparation of DNA for Pulsed-field Gel Electrophoresis: Isolation of DNA from<br>Mammalian Cells and Tissues         | 5.61     |
|                              | 14 Preparation of DNA for Pulsed-field Gel Electrophoresis: Isolation of Intact DNA from Yeast                           | 5.65     |
|                              | 15 Restriction Endonuclease Digestion of DNA in Agarose Plugs  | 5.68     |
|                              | 16 Markers for Pulsed-field Gel Electrophoresis  | 5,71     |
|                              | 17 Pulsed-field Gel Electrophoresis via Transverse Alternating Field Electrophoresis Gels                                | 5.74     |
|                              | Alternative Protocol: Silver Staining PFGE Gels  | 5.77     |
|                              | 18 Pulsed-field Gel Electrophoresis via Contour-clamped Homogeneous Electric Field G                                     | els 5,79 |
|                              | 19 Direct Retrieval of DNA Fragments from Pulsed-field Gels  | 5.83     |
|                              | 20 Retrieval of DNA Fragments from Pulsed-field Gels following DNA Concentration   | 5.86     |

#### Contents

| Chapter 6   |            |            |             |          |
|-------------|------------|------------|-------------|----------|
| Preparation | and Analy: | sis of Euk | aryotic Ger | omic DNA |

6.1

### INTRODUCTION PROTOCOLS

| ation of High-molecular-weight DNA from Mammalian Cells Using Proteinase K Phenol diditional Protocol: Estimating the Concentration of DNA by Fluorometry ation of High-molecular-weight DNA from Mammalian Cells Using Formamide ation of DNA from Mammalian Cells by Spoofing ation of DNA from Mammalian Cells Grown in 96-well Microtiter Plates Iditional Protocol: Optimizing Genomic DNA Isolation for PCR haration of Genomic DNA from Mammalian Cells around the Small Samples | 6.4<br>6.12<br>6.13<br>6.16<br>6.19<br>6.22  |
|---|--|
| ation of High-molecular-weight DNA from Mammalian Cells Using Formamide \ ation of DNA from Mammalian Cells by Spooling ation of DNA from Mammalian Cells Grown in 96-well Microtiter Plates iditional Protocol: Optimizing Genomic DNA Isolation for PCR arration of Genomic DNA from Mouse Talls and Other Small Samples  | 6.13<br>6.16<br>6.19<br>6.22   |
| ation of DNA from Mammalian Cells by Spooling<br>ation of DNA from Mammalian Cells Grown in 96-well Microtiter Plates<br>iditional Protocol: Optimizing Genomic DNA Isolation for PCR<br>aration of Genomic DNA from Mouse Talls and Other Small Samples  | 6.16<br>6.19<br>6.22   |
| ation of DNA from Mammalian Cells by Spooling<br>ation of DNA from Mammalian Cells Grown in 96-well Microtiter Plates<br>iditional Protocol: Optimizing Genomic DNA Isolation for PCR<br>aration of Genomic DNA from Mouse Talls and Other Small Samples  | 6.19<br>6.22   |
| ation of DNA from Mammalian Cells Grown in 96 well Microtiter Plates<br>Iditional Protocol: Optimizing Genomic DNA Isolation for PCR<br>paration of Genomic DNA from Mouse Tails and Other Small Samples  | 6.22   |
| Iditional Protocol: Optimizing Genomic DNA Isolation for PCR<br>paration of Genomic DNA from Mouse Talls and Other Small Samples  |  |
| paration of Genomic DNA from Mouse Tails and Other Small Samples  |  |
|   | 6.23   |
| ternative Protocol: Isolation of DNA from Mouse Tails without Extraction by ganic Solvents  | 6.26   |
| ternative Protocol: One-tube Isolation of DNA from Mouse Tails  | 6.26   |
| ternative Protocol: DNA Extraction from Paraffin Blocks   | 6.27   |
| d Isolation of Mammalian DNA  | 6.28   |
| d Isolation of Yeast DNA  | 6.31   |
| ction to Southern Hybridization (Protocols 8-10)  | 6.33   |
| thern Blotting: Capillary Transfer of DNA to Membranes  | 6.39   |
| thern Blotting: Simultaneous Transfer of DNA from a Single Agarose Gel to<br>Membranes  | 6.47   |
| thern Hybridization of Radiolabeled Probes to Nucleic Acids Immobilized on<br>obranes   | 6.50   |
| ditional Protocol: Stripping Probes from Membranes  | 6.57   |
| ditional Protocol: Hybridization at Low Stringency  | 6.58   |
|   | hern Blotting: Simultaneous Transfer of DNA from a Single Agarose Gel to<br>Membranes<br>hern Hybridization of Radiolabeled Probes to Nucleic Acids Immobilized on<br>branes<br>ditional Protocol: Stripping Probes from Membranes |

## Chapter 7

| Extraction, | Purification, | and | Analysis | of mRNA from | n |
|-------------|---------------|-----|----------|--------------|---|
| Eukaryotic  | Cells         |     |          |              |   |

Rapid Hybridization Buffers

Formamide and Its Uses in Molecular Cloning

Spooling DNA (Historical Footnote)

7.1

6.59

6.61

6.61

6.62

### INTRODUCTION

CTAB

### PROTOCOLS

| 1 | Purification of RNA from Cells and Tissues by Acid Phenol-Guanidinium          | 7.4 |
|---|--|-----|
|   | Thiocyanate-Chloroform Extraction  |     |
| 2 | A Single-step Method for the Simultaneous Preparation of DNA, RNA, and Protein | 7.9 |

from Cells and Tissues

Contents хi

| ,                    | Selection of Poly(A)* RNA by Oligo(dT)-Cellulose Chromatography                       | 7.13 |
|----------------------|---|------|
|                      | Selection of Poly(A)+ RNA by Batch Chromatography                                     | 7.18 |
|                      | troduction to Northern Hybridization (Protocols 5–9)                                  | 7.21 |
|                      | Separation of RNA According to Size: Electrophoresis of Clyoxylated RNA through       | 7.27 |
| ,                    | Agarose Gels  |      |
|                      | Separation of RNA According to Size: Electrophoresis of RNA through Agarose Gels      | 7.31 |
| 0                    | Containing formaldehyde   |      |
| 7                    | Transfer and Fixation of Denatured RNA to Membranes                                   | 7.35 |
| •                    | Alternative Protocol: Capillary Transfer by Downward Flow                             | 7.41 |
| 8                    | Northern Hybridization  | 7.42 |
|                      | Dot and Slot Hybridization of Purified RNA  | 7.46 |
|                      | Mapping RNA with Nuclease S1  | 7.51 |
|                      | Ribonuclease Protection: Mapping RNA with Ribonuclease and Radiolabeled RNA           | 7.63 |
|                      | Probes  |      |
| 12                   | Analysis of RNA by Primer Extension   | 7.75 |
|                      |   |      |
| INFOR                | RMATION PANELS  |      |
| Ho                   | w to Win the Battle with RNase  | 7.82 |
| Inh                  | ibitors of RNases   | 7.83 |
| Die                  | ethylpyrocarbonate  | 7.84 |
| Gu                   | anidinium Salts   | 7.85 |
| Nu                   | clease S1   | 7.86 |
| Exc                  | onuclease VII   | 7.86 |
|                      | ing Bean Nuclease   | 7.87 |
| Pro                  | moter Sequences Recognized by Bacteriophage-encoded RNA Polymerases                   | 7.87 |
| Act                  | tinomycin D   | 7.88 |
| INDEX                | ÇI.T  |      |
|                      | Volume 2  |      |
| Chapter 8            |   |      |
| Chapter 8            |   | 8.1  |
| In Vitro Amplificati | ion of DNA by the Polymerase  | 0.1  |
| Chain Reaction       |   |      |
| IN IED C             | DUCTION:  |      |
|                      | DUCTION   |      |
| PROTO                | ocols   |      |
| 1                    | The Basic Polymerase Chain Reaction   | 8.18 |
| 2                    |   | 8.25 |
| 3                    | Removal of Oligonucleotides and Excess dNTPs from Amplified DNA by Ultrafiltration    | 8.27 |
| In                   | troduction to Cloning PCR Products (Protocols 4–7)                                    | 8.30 |
| 4                    | Blunt-end Cloning of PCR Products   | 8.32 |
|                      | Cloning PCR Products into T Vectors   | 8.35 |
| 6                    | Cloning PCR Products by Addition of Restriction Sites to the Termini of Amplified DNA | 8.37 |
| 7                    |   | 8.42 |
| 8                    | Amplification of cDNA Generated by Reverse Transcription of mRNA (RT-PCR)             | 8.46 |
|                      |   |      |

| xii | Contents |
|-----|----------|

| 9 Rapid Amplification of 5" cDNA Ends (5"-RACE)   | 8.54    |
|---|---------|
| 10 Rapid Amplification of 3 CDNA Ends (3 -RACE)   | 8.61    |
| 11 Mixed Oligonucleotide-primed Amplification of cDNA (MOPAC)   | 8.66    |
| 12 Rapid Characterization of DNAs Cloned in Prokaryotic Vectors   | 8.72    |
| <ul> <li>Additional Protocol: Screening Yeast Colonies by PCR</li> </ul>  | 8.75    |
| <ul> <li>Additional Protocol: Screening Bacteriophage λ Libraries</li> </ul>  | 8.76    |
| 13 Long PCR   | 8.77    |
| 14 Inverse PCR  | 8.81    |
| 15 Quantitative PCR   | 8.86    |
| 16 Differential Display–PCR (DD-PCR)  | 8.96    |
| INFORMATION PANELS  |         |
| Multiplex PCR   | 8.107   |
| Taq DNA Polymerase  | 8.108   |
| Hot Start PCR   | 8,110   |
| Ribonuclease H  | 8,111   |
| Terminal Transferase  | 8.111 . |
| Touchdown PCR Use of Inosine in Degenerate Pools of Oligonucleotides Used for PCR   | 8.113   |
| Universal Primers   | 8.113   |
|   |         |
| Chapter 9   |         |
| Preparation of Radiolabeled DNA and RNA Probes  | 9.1     |
| INTRODUCTION  |         |
| PROTOCOLS   |         |
| 1 Random Priming: Radiolabeling of Purified DNA Fragments by Extension of Random  | 9.4     |
| Oligonucleotides  | 2.4     |
| 2 Random Priming: Radiolabeling of DNA by Extension of Random Oligonucleotides in   | 9.9     |
| the Presence of Melted Agarose  |         |
| <ul> <li>Nick Translation: An Historical Note</li> </ul>  | 9.12    |
| 3 Radiolabeling of DNA Probes by the Polymerase Chain Reaction  | 9.14    |
| <ul> <li>Additional Protocol: Asymmetric Probes</li> </ul>  | 9.18    |
| 4 Synthesis of Single-stranded DNA Probes of Defined Length from Bacteriophage  | 9.19    |
| M13 Templates   |         |
| 5 Synthesis of Single-stranded DNA Probes of Heterogeneous Length from<br>Bacteriophage M13 Templates   | 9.25    |
| 6 Synthesis of Single-stranded RNA Probes by In Vitro Transcription   | 9.29    |
| Additional Protocol: Using PCR to Add Promoters for Bacteriophage-encoded   | 9.36    |
| RNA Polymerases to Fragments of DNA   | 3,30    |
| 7 Synthesis of cDNA Probes from mRNA Using Random Oligonucleotide Primers   | 9.38    |
| 8 Synthesis of Radiolabeled, Subtracted cDNA Probes Using Oligo(dT) as a Primer   | 9.41    |
| 9 Radiolabeling of Subtracted cDNA Probes by Random Oligonucleotide Extension   | 9.46    |
| 10 Labeling 3" Termini of Double-stranded DNA Using the Klenow Fragment of E. coli  | 9.51    |
| DNA Polymerase I  |         |
| 11 Labeling 3' Termini of Double-stranded DNA with Bacteriophage T4 DNA   | 9.57    |
| Polymerase  |         |
| 12 End Labeling Protruding 3' Termini of Double-stranded DNA with (σ- <sup>22</sup> P)Cordycepin<br>5' Triphosphate or (α- <sup>32</sup> P)Idideoxy ATP | 9.60    |

|                    | Content   | s xiii         |
|--------------------|---|----------------|
| 13                 | Dephosphorylation of ONA Fragments with Alkaline Phosphatase  | 9.62           |
| 14                 |   | 9.66           |
|                    | Phosphorylation of DNA Molecules with Dephosphorylated Blunt Ends or Recessed 5* Termini  |                |
| 16                 | Phosphorylation of DNA Molecules with Protruding 5° Termini by the Exchange Reaction  | 9.73           |
| INFO               | RMATION PANELS  |                |
| 1                  | Nonradioactive Labeling of Nucleic Acids  | 9.76           |
| E                  | E. coli DNA Polymerase I and the Klenow Fragment  | 9.82           |
|                    | In Vitro Transcription Systems  | 9.87           |
|                    | solating Differentially Expressed cDNAs by Differential Screening and Cloning<br>Alkaline Phosphatase                           | 9.89<br>9.92   |
| Chapter 10         |   |                |
| Working with Synt  | thetic Oligonucleotide Probes   | 10.1           |
| INTRO              | DDUCTION  |                |
|                    | DCOLS   |                |
|                    |   | 10.11          |
| 1                  | Purification of Synthetic Oligonucleotides by Polyacrylamide Gel Electrophoresis  | 10.11<br>10.17 |
| 3                  | Phosphorylating the 5´ Termini of Oligonucleotides  Purification of Radiolabeled Oligonucleotides by Precipitation with Ethanol | 10.17          |
| 4                  |   | 10.22          |
| 5                  | Purification of Radiolabeled Oligonucleotides by Size-exclusion Chromatography  | 10.25          |
| 6                  | Purification of Radiolabeled Oligonucleotides by Chromatography on a Sep-Pak $C_{1\delta}$ Column                               | 10.28          |
|                    | Labeling of Synthetic Oligonucleotides Using the Klenow Fragment of £. coli DNA Polymerase I                                    | 10.30          |
| 8                  | Hybridization of Oligonucleotide Probes in Aqueous Solutions: Washing in Buffers Containing Quaternary Ammonium Salts           | 10.35          |
| , 9                | Empirical Measurement of Melting Temperature  | 10.38          |
| INFOR              | MATION PANELS   |                |
|                    | zonucleotide Synthesis  | 10.42          |
|                    | ting Temperatures<br>thods Used to Purify Synthetic Oligonucleotides  | 10.47<br>10.48 |
| Chapter 11         |   |                |
| Preparation of cDN | NA Libraries and Gene Identification  | 11.1           |
| INTRO              | DUCTION   |                |
| PROTO              | OCOLS   |                |
| 1                  | Construction of cDNA Libraries  | 11.38          |
|                    | Stage 1: Synthesis of First-strand cDNA Catalyzed by Reverse Transcriptase  | 11.39          |
|                    | Stage 2: Second-strand Synthesis  | 11.44          |

11.49

12.31

12.32

#### xiv Contents

|              | Stage 3: Methylation of cDNA  | 11.49  |
|--------------|---|--------|
|              | Stage 4: Attachment of Linkers or Adaptors  | 11.52  |
|              | Stage 5: Fractionation of cDNA by Gel Filtration through Sephanose CL-4B  | 11.57  |
|              | Stage 6: Ligation of cDNA to Bacteriophage \(\lambda\) Arms   | 11.60  |
|              | Alternative Protocol: Ligation of cDNA into a Plasmid Vector  | 11.64  |
|              | <ul> <li>Additional Protocol: Amplification of cDNA Libraries</li> </ul>  | 11.65  |
|              | 2 Construction and Screening of Eukaryotic Expression Libraries   | 11.68  |
|              | Stage 1: Construction of cDNA Libraries in Eukaryotic Expression Vectors  | 11.69  |
|              | Stage 2: Screening cDNA Libraries Constructed in Eukaryotic Expression Vectors  | 11.75  |
|              | 3 Exon Trapping and Amplification   | 11.80  |
|              | Stage 1: Construction of the Library  | 11.82  |
|              | Stage 2: Electroporation of the Library into COS-7 Cells  | 11.87  |
|              | Stage 3: Harvesting the mRNA  | 11.89  |
|              | Stage 4: Reverse Transcriptase-PCR  | 11.91  |
|              | Stage 5: Analysis of Clones   | 11.96  |
|              | 4 Direct Selection of cDNAs with Large Genomic DNA Clones   | 11.98  |
|              | INFORMATION PANELS  |        |
|              | Commercial Kits for cDNA Synthesis and Library Construction   | 11.109 |
|              | Mo-MLV Reverse Transcriptase  | 11.111 |
|              | Homopolymeric Tailing   | 11.112 |
|              | Agt10 and Agt11   | 11.113 |
|              | Constructing cDNA Libraries from Small Numbers of Cells   | 11.114 |
|              | In Vitro Packaging  | 11,115 |
|              | COS Cells   | 11.116 |
|              | Biotin  | 11.117 |
|              | Magnetic Beads  | 11.120 |
|              | Ligation-independent Cloning  | 11.123 |
| Chapter 12   |   |        |
| DNA Sequence | ing   | 12.1   |
|              | INTRODUCTION  |        |
|              | PROTOCOLS   |        |
|              | Generation of a Library of Randomly Overlapping DNA Inserts   | 12.10  |
|              | Alternative Protocol: Preparation of Small Numbers of Single-stranded DNA   | 12.10  |
|              | Templates from Bacteriophage M13  |        |
|              | <ul> <li>Additional Protocol: Preparation of Dephosphorylated Blunt-ended Bacteriophage<br/>M13 Vector DNA for Shotgun Cloning</li> </ul> | 12.24  |
|              | 2 Preparing Denatured Templates for Sequencing by Dideoxy-mediated Chain<br>Termination   | 12.26  |
|              | - Additional Protocol: Paoid Denaturation of Double-stranded DNA  | 12.30  |

• Additional Protocol: Purification of Plasmid DNA from Small-scale Cultures by

3 Dideoxy-mediated Sequencing Reactions Using Bacteriophage T7 DNA Polymerase

Precipitation with PEG

(Sequenase)

|             |       | · · · · · · · · · · · · · · · · · · ·   | -      |                  |
|-------------|-------|---|--------|------------------|
|             |       |   |        |                  |
|             | 4     | Dideoxy-mediated Sequencing Reactions Using the Klenow Fragment of E. co.<br>DNA Polymerase I and Single-stranded DNA Templates                             | li     | 12.40            |
|             | 5     | Dideoxy-mediated Sequencing of DNA Using Taq DNA Polymerase   |        | 12.45            |
|             | 6     | Cycle Sequencing: Dideoxy-mediated Sequencing Reactions Using PCR and   |        | 12.51            |
|             |       | End-labeled Primers   |        |                  |
|             |       | <ul> <li>Additional Protocol: Cycle Sequencing Reactions Using PCR and Internal</li> </ul>  |        | 12.60            |
|             |       | Labeling with [a-32P]dNTPs  |        | 12.61            |
|             | 7     | Chemical Sequencing   |        | 12.70            |
|             |       | <ul> <li>Alternative Protocol: Rapid Maxam-Gilbert Sequencing</li> <li>Additional Protocol: Preparation of End-labeled DNA for Chemical Sequence</li> </ul> | ine    | 12.73            |
|             |       | Preparation of Denaturing Polyacrylamide Gels   |        | 12.74            |
|             |       | Preparation of Denaturing Polyacrylamide Gels Containing Formamide  |        | 12.81            |
|             |       | Preparation of Electrolyte Gradient Gels  |        | 12.83            |
|             |       | Loading and Running DNA-sequencing Gels   |        | 12.85            |
|             | 12    |   |        | 12.90            |
|             |       |   |        |                  |
|             | INFOR | RMATION PANELS  |        |                  |
|             | A     | outomated DNA Sequencing  |        | 12.94            |
|             | ٨     | Aicrotiter Plates   |        | 12.100           |
|             | т     | he Klenow Fragment of E. coli DNA Polymerase 1  |        | 12,101           |
|             | P     | Preparation of Stock Solutions of Oligonucleotide Primers for DNA Sequencing  |        | 12.103           |
|             |       | equenase  |        | 12.104<br>12.106 |
|             | C     | Conventional Chain-termination Sequencing of PCR-amplified DNA  |        | 12.106           |
|             |       | Preparation of Stock Solutions of dNTPs and ddNTPs for DNA Sequencing   |        | 12.107           |
|             |       | Clycerol in DNA Sequencing Reactions  |        | 12.109           |
|             |       | Compressions in DNA Sequencing Gels   |        | 12.111           |
|             |       | r-deaza-dGTP  |        | 12.112           |
|             |       | Dichlorodimethylsilane<br>Reading an Autoradiograph   |        | 12.113           |
|             |       | lectrical Mobility of DNA   |        | 12,114           |
|             | ٠     | nectical Property of Divi   |        |                  |
|             |       |   |        |                  |
| Chapter 13  |       |   |        | 101              |
| Mutagenesis |       |   |        | 13.1             |
|             | INTRO | ODUCTION .  |        |                  |
|             |       | ocols   |        |                  |
|             |       | Preparation of Uracil-containing Single-stranded Bacteriophage M13 DNA  |        | 13.11            |
|             | 2     | Oligonucleotide-directed Mutagenesis of Single-stranded DNA   |        | 13.15            |
|             | 3     | In Vitro Mutagenesis Using Double-stranded DNA Templates: Selection of Mi   | utants | 13.19            |
|             |       | with Dpnl   |        |                  |
|             | 4     | Oligonucleotide-directed Mutagenesis by Elimination of a Unique Restriction (USE Mutagenesis)   | Site   | 13.26            |
|             | 5     | Rapid and Efficient Site-directed Mutagenesis by the Single-tube Megaprimer<br>Method   | PCR    | 13.31            |
|             | 6     | Site-specific Mutagenesis by Overlap Extension  |        | 13.36            |
|             |       |   |        |                  |

### xvi Contents

Chapter 14 Screening

|    | 7  | Screening Recombinant Clones for Site-directed Mutagenesis by Hybridization to<br>Radiolabeled Oligonucleotides  | 13.40  |
|----|--|--|--|
|    |  | <ul> <li>Alternative Protocol: Screening Phagemid-containing Bacterial Colonies by</li> </ul>  | 13.47  |
|    |  | Hybridization to Radiolabeled Oligonucleotides   | 13.48  |
|    |  | Alternative Protocol: Detection of Defined Mutants by PCR  | 13.49  |
|    | 8  | Detection of Mutations by Single-strand Conformational Polymorphism and Heteroduplex Analysis  | 13,43  |
|    | 9  | Generation of Sets of Nested Deletion Mutants with Exonuclease III   | 13.57  |
|    | 10   | Generation of Bidirectional Sets of Deletion Mutants by Digestion with BAL 31 Nuclease   | 13.62  |
|    | INFOR  | MATION PANELS  |  |
|    | ε  | AL 31  | 13.68  |
|    | 6  | xonuclease III   | 13.72  |
|    | i.   | inker-scanning Mutagenesis   | 13.75 \  |
|    | F  | andom Mutagenesis  | 13.78  |
|    | A  | lanine-scanning Mutagenesis  | 13.81  |
|    | N  | Autagenic Oligonucleotides   | 13.82  |
|    | S  | electing against Wild-type DNA in Site-directed Mutagenesis  | 13.84  |
|    | ^  | 6-methyladenine, Dam Methylase, and Methylation-sensitive Restriction Enzymes  | 13.87  |
|    | C  | ommercial Kits for Site-directed Mutagenesis   | 13.89  |
|    | C  | lycerol  | 13.90  |
|    | ٨  | futation Dete⊂tion .   | 13.91  |
|    |  |  |  |
| Εx | pressi   | on Libraries   | 14.1   |
| E× |  | on Libraries   | 14.1   |
| E× |  | DUCTION  | 14.1   |
| Ex | INTRO  | DUCTION  | 14.1   |
| Ex | INTRO<br>PROTO   | DUCTION  |  |
| Ex | INTRO PROTO  | DUCTION DCOLS  Screening Expression Libraries Constructed in Bacteriophage \( \) Vectors  Screening Expression Libraries Constructed in Plasmid Vectors  Removal of Cross-reactive Antibodies from Antiserum: Pseudoscreening  | 14.4   |
| Ex | INTRO PROTO 1 2  | DUCTION  COLS  Screening Expression Libraries Constructed in Bacteriophage λ Vectors Screening Expression Libraries Constructed in Plasmid Vectors Removal of Cross-reactive Antibodies from Antiserum: Pseudoscreening  • Alternative Protocol: Adsorbing Antibodies with Lysates of Bacteriophage-   | 14.4   |
| Ex | PROTO 1 2 3  | DUCTION DOCIS  Screening Expression Libraries Constructed in Bacteriophage \( \lambda \) Vectors Screening Expression Libraries Constructed in Plasmid Vectors Removal of Cross-reactive Antibodies from Antiserum: Pseudoscreening  • Alternative Protocol: Adsorbing Antibodies with Lysates of Bacteriophage- infected Cells  | 14.4<br>14.14<br>14.23<br>14.25  |
| Ex | INTRO PROTO 1 2 3                                      | DUCTION  COLS  Screening Expression Libraries Constructed in Bacteriophage λ Vectors  Screening Expression Libraries Constructed in Plasmid Vectors  Removal of Cross-reactive Antibodies from Antiserum: Peacuboscreening  • Alternative Protocol: Adsorbing Antibodies with Lysates of Bacteriophage- infected Cells  Removal of Cross-reactive Antibodies from Antiserum: Incubation with ε. coll Lysate  | 14.4<br>14.14<br>14.23<br>14.25  |
| Ex | INTRO PROTO 1 2 3                                      | DUCTION  DOCIS  Screening Expression Libraries Constructed in Bacteriophage λ Vectors Screening Expression Libraries Constructed in Plasmid Vectors Removal of Cross-reactive Antibodies from Antiserum: Pseudoscreening  • Alternative Protocol: Adsorbing Antibodies with Lysates of Bacteriophage- infected Cells Removal of Cross-reactive Antibodies from Antiserum: Incubation with £, coli Lysate Removal of Cross-reactive Antibodies from Antiserum: Affinity Chromatography  | 14.4<br>14.14<br>14.23<br>14.25  |
| Ex | INTRO PROTO 1 2 3                                      | DUCTION  COLS  Screening Expression Libraries Constructed in Bacteriophage λ Vectors  Screening Expression Libraries Constructed in Plasmid Vectors  Removal of Cross-reactive Antibodies from Antiserum: Pseudoscreening  • Alternative Protocol: Adsorbing Antibodies with Lysates of Bacteriophage- infected Cells  Removal of Cross-reactive Antibodies from Antiserum: Incubation with ε. coli Lysate  Removal of Cross-reactive Antibodies from Antiserum: Affinity Chromatography  Identifying DNA-binding Proteins in Bacteriophage λ Expression Libraries  Preparation of Lysates Containing Fusion Proteins Encoded by Bacteriophage λ   | 14.4<br>14.14<br>14.23<br>14.25<br>14.26<br>14.28                            |
| Ex | INTRO<br>PROTO<br>1<br>2<br>3<br>4<br>5<br>6<br>7      | DUCTION  COLS  Screening Expression Libraries Constructed in Bacteriophage λ Vectors  Screening Expression Libraries Constructed in Plasmid Vectors  Removal of Cross-reactive Antibodies from Antiserum: Pecudoscreening  • Alternative Protocol: Adsorbing Antibodies with Lysates of Bacteriophage- infected Cells  Removal of Cross-reactive Antibodies from Antiserum: Incubation with ε. coll Lysate  Removal of Cross-reactive Antibodies from Antiserum: Affinity Chromatography  Identifying DN-Abinding Protonis in Bacteriophage a Expression Libraries  Preparation of Lysates Containing Fusion Proteins Encoded by Bacteriophage λ  Lysogens: Lysis of Bacterial Colonies  | 14.4<br>14.14<br>14.23<br>14.25<br>14.26<br>14.28<br>14.31                   |
| Ex | INTRO PROTC  1 2 3 4 5 6 7 8                           | DUCTION  COLS  Screening Expression Libraries Constructed in Bacteriophage λ Vectors Screening Expression Libraries Constructed in Plasmid Vectors Removal of Cross-reactive Antibodies from Antiserum: Pseudoscreening  • Alternative Protocol: Adsorbing Antibodies with Lysates of Bacteriophage- infected Cells Removal of Cross-reactive Antibodies from Antiserum: Incubation with ε, coli Lysate Removal of Cross-reactive Antibodies from Antiserum: Affinity Chromatography Identifying DNA-binding Proteins in Bacteriophage & Expression Libraries Preparation of Lysates Containing Fusion Proteins Encoded by Bacteriophage λ Lysogens: Lysis of Bacterial Colonies Preparation of Lysates Containing fusion Proteins Encoded by Bacteriophage λ: Lytic Infections on Agar Plates   | 14.4<br>14.14<br>14.23<br>14.25<br>14.26<br>14.28<br>14.31<br>14.37          |
| Ex | INTRO<br>PROTO<br>1<br>2<br>3<br>4<br>5<br>6<br>7      | DUCTION  COLS  Screening Expression Libraries Constructed in Bacteriophage λ Vectors  Screening Expression Libraries Constructed in Plasmid Vectors  Removal of Cross-reactive Antibodies from Antiserum: Pseudoscreening  • Alternative Protocol: Adsorbing Antibodies with Lysates of Bacteriophage- infected Cells  Removal of Cross-reactive Antibodies from Antiserum: Incubation with ε. coll Lysate  Removal of Cross-reactive Antibodies from Antiserum: Affinity Chromatography  Identifying DNA-binding Proteins in Bacteriophage & Expression Libraries  Preparation of Lysates Containing Fusion Proteins Encoded by Bacteriophage λ  Lysogens: Lysis of Bacterial Colonies  Preparation of Lysates Containing Fusion Proteins Encoded by Bacteriophage λ:  Lytic Infections on Agar Plates  | 14.4<br>14.14<br>14.23<br>14.25<br>14.26<br>14.28<br>14.31                   |
| Ex | INTRO<br>PROTO<br>1<br>2<br>3<br>4<br>5<br>6<br>7<br>8 | DUCTION  DOCIS  Screening Expression Libraries Constructed in Bacteriophage λ Vectors  Screening Expression Libraries Constructed in Plasmid Vectors  Removal of Cross-reactive Antibodies from Antiserum: Pseudoscreening  • Alternative Protocol: Adsorbing Antibodies with Lysates of Bacteriophage- infected Cells  Removal of Cross-reactive Antibodies from Antiserum: Incubation with ε. coll Lysate  Removal of Cross-reactive Antibodies from Antiserum: Affinity Chromatography  Identifying DNA-binding Proteins in Bacteriophage ε. Expression Libraries  Preparation of Lysates Containing Fusion Proteins Encoded by Bacteriophage λ  Lystic Infections on Agar Plates  Preparation of Lysates Containing Fusion Proteins Encoded by Bacteriophage λ:  Lytic Infections on Agar Plates   | 14.4<br>14.14<br>14.23<br>14.25<br>14.26<br>14.28<br>14.31<br>14.37          |
| Ex | INTRO PROTO 1 2 3 4 5 6 6 7 8 9 INFOR                  | DUCTION  COLS  Screening Expression Libraries Constructed in Bacteriophage λ Vectors  Screening Expression Libraries Constructed in Plasmid Vectors  Screening Expression Libraries Constructed in Plasmid Vectors  Removal of Cross-reactive Antibodies from Antiserum: Eucodoscreening  * Alternative Protocol: Adsorbing Antibodies with Lysates of Bacteriophage- infected Cells  Removal of Cross-reactive Antibodies from Antiserum: Incubation with Ε. coll Lysate  Removal of Cross-reactive Antibodies from Antiserum: Affinity Chromatography  Identifying DN-binding Protonis in Bacteriophage & Expression Libraries  Preparation of Lysates Containing Fusion Proteins Encoded by Bacteriophage λ  Lysic Infections on Agar Plates  Preparation of Lysates Containing Fusion Proteins Encoded by Bacteriophage λ:  Lytic Infections in Liquid Medium  MATION PANELS                                 | 14.4<br>14.14<br>14.23<br>14.25<br>14.26<br>14.28<br>14.31<br>14.37          |
| Ex | INTRO PROTO 1 2 3 4 5 6 7 8 9 INFOR                    | DUCTION  DICOLS  Screening Expression Libraries Constructed in Bacteriophage λ Vectors Screening Expression Libraries Constructed in Plasmid Vectors Removal of Cross-reactive Antibodies from Antiserum: Pseudoscreening  • Alternative Protocol: Adsorbing Antibodies with Lysates of Bacteriophage- infected Cells Removal of Cross-reactive Antibodies from Antiserum: Incubation with ε, coli Lysate Removal of Cross-reactive Antibodies from Antiserum: Affinity Chromatography Identifying DNA-binding Proteins in Bacteriophage & Expression Libraries Preparation of Lysates Containing Fusion Proteins Encoded by Bacteriophage λ Lysogens: Lysis of Bacterial Colonies Preparation of Lysates Containing Fusion Proteins Encoded by Bacteriophage λ; Lytic Infections on Agar Plates Preparation of Lysates Containing Fusion Proteins Encoded by Bacteriophage λ; Lytic Infections in Liquid Medium | 14.4<br>14.14<br>14.23<br>14.25<br>14.26<br>14.28<br>14.31<br>14.37<br>14.41 |

Contents xvii

| Volume 3   |   |
|--|---|
| Chapter 15   |   |
| Expression of Cloned Genes in Escherichia coli   | 15.1  |
| INTRODUCTION PROTOCOLS   |   |
| 1 Expression of Cloned Genes in £. coli Using IPTG-inducible Promoters 2 Expression of Cloned Genes in £. coli Using the Bacteriophage Τ? Promoter 3 Expression of Cloned Genes in £. coli Using the Bacteriophage λ. ρ. Promoter 4 Expression of Secreted Foreign Proteins Using the Alkaline Phosphatase Promoter (ρ)πολ) and Signal Sequence • Additional Protocol: Subcellular Localization of PhoA Fusion Proteins 5 Purification of Fusion Proteins by Affinity Chromatography on Glutathione Agarose 6 Purification of Maltose-binding Fusion Proteins by Affinity Chromatography on Amylose Resin 7 Purification of Histidine-tagged Proteins by Immobilized Ni²+ Absorption Chromatography • Alternative Protocol: Elution of Polyhistidine-tagged Proteins from Metal Affinity Columns Using Decreasing pH • Additional Protocol: Regeneration of NTA-Ni²+-Agarose 8 Purification of Expressed Proteins from IntA-Ni²- | 15.14<br>15,20<br>15.25<br>15.30<br>15.35<br>15.36<br>15.40<br>15,44<br>15,47 |
| <ul> <li>Additional Protocol: Refolding Solubilized Proteins Recovered from Inclusion<br/>Bodies</li> </ul>  | 15.53   |
| INFORMATION PANELS   |   |
| Expression of Cloned Genes   | 15.55   |
| E. coli Expression Systems   | 15.56   |
| LacZ Fusions<br>Chaotropic Agents  | 15.57<br>15.60  |
| Chapter 16   | 16.1  |
| Introducing Cloned Genes into Cultured Mammalian Cells   | 10.1  |
| INTRODUCTION PROTOCOLS   |   |
| <ol> <li>DNA Transfection Mediated by Lipofection</li> <li>Additional Protocol: Histochemical Staining of Cell Monolayers for<br/>B-Galactosidase</li> <li>Calcium-phosphate-mediated Transfection of Eukaryotic Cells with Plasmid DNAs</li> </ol>  | 16.7<br>16.13<br>16.14  |
| <ul> <li>Alternative Protocol: High-efficiency Calcium-phosphate-mediated Transfection<br/>of Eukaryotic Cells with Plasmid DNAs</li> </ul>  | 16.19   |
| <ol> <li>Calcium-phosphate-mediated Transfection of Cells with High-molecular-weight<br/>Genomic DNA</li> </ol>  | 16.21   |

#### xviii Contents

|       | <ul> <li>Alternative Protocol: Calcium-phosphate-mediated Transfection of Adherent<br/>Cells</li> </ul> | 16.25 |
|-------|---|-------|
|       | Alternative Protocol: Calcium-phosphate-mediated Transfection of Cells<br>Growing in Suspension         | 16.26 |
| 4     | Transfection Mediated by DEAE-Dextran: High-efficiency Method   | 16.27 |
|       | Alternative Protocol: Transfection Mediated by DEAE-Dextran: Increased Cell Viability                   | 16.32 |
| 5     | DNA Transfection by Electroporation   | 16.33 |
| 6     | DNA Transfection by Biolistics  | 16.37 |
|       | Additional Protocol: Histochemical Staining of Cell Monolayers or     Tissue for β-Glucuronidase        | 16.42 |
| 7     | DNA Transfection Using Polybrene  | 16.43 |
| INFOR | MATION PANELS   | /     |
|       | Cotransformation  | 16.47 |
|       | Sciective Agents for Stable Transformation  | 16.48 |
|       | Lipofection   | 16.50 |
|       | Transfection of Mammalian Cells with Calcium Phosphate-DNA Coprecipitates                               | 16.52 |
|       | Chloroquine Diphosphate   | 16.53 |
|       | Electroporation   | 16.54 |

### Chapter 17

| Chapter 17  |       |
|---|-------|
| Analysis of Gene Expression in Cultured Mammalian C | Cells |
|   |       |

Using Thin-layer Chromatography

Products into Scintillation Fluid
6 Assay for Luciferase in Extracts of Mammalian Cells

17.1

17,40

17.41

17.42

## INTRODUCTION PROTOCOLS

### Cis-acting Regions and Trans-acting Factors

|    | Delta by Obland I Spotagining  | 17.4  |
|----|--|-------|
| 1  | Mapping Protein-binding Sites on DNA by DNase I Footprinting                               |       |
|    | <ul> <li>Alternative Protocol: Mapping Protein-binding Sites on DNA by Hydroxyl</li> </ul> | 17.12 |
|    | Radical Footprinting   |       |
| 2  | Gel Retardation Assays for DNA-binding Proteins  | 17.13 |
| •  |  | 17,17 |
|    | Additional Protocol: Supershift Assays   |       |
|    | Additional Protocol: Competition Assays  | 17.17 |
| 3  | Mapping DNase-I-hypersensitive Sites   | 17,18 |
| Α  | nalysis of Primary Transcripts   |       |
| 4  | Transcriptional Run-on Assays  | 17,23 |
| R  | eporter Assays   |       |
| 1r | stroduction to Reporter Assays: CAT, Luciferase, and β-galactosidase (Protocols 5-7)       | 17.30 |
| 5  | Measurement of Chloramphenicol Acetyltransferase in Extracts of Mammalian Cells            | 17.33 |

Alternative Protocol: Measurement of CAT by Extraction with Organic Solvents

· Alternative Protocol: Measurement of CAT following Diffusion of Reaction

3

|            | Conten   | ts xix         |
|------------|--|----------------|
|            | Alternative Protocol: Using a Scintillation Counter to Measure Luciferase     Alternative Protocol: Using a Scintillation Counter to Measure Luciferase                | 17.46<br>17.47 |
|            | <ul> <li>Alternative Protocol: Assay for Luciferase in Cells Growing in 96-well Plates</li> <li>Assay for β-galactosidase in Extracts of Mammalian Cells</li> </ul>    | 17.48          |
|            | Inducible Systems  |                |
|            | 8 Tetracycline as Regulator of Inducible Gene Expression in Mammalian Cells  | 17.52          |
|            | Stage 1: Stable Transfection of Fibroblasts with pTet-tTAk   | 17.60          |
|            | Stage 2: Stable Transfection of Inducible tTA-expressing NIH-3T3 Cells with<br>Tetracycline-regulated Target Genes   | 17.65          |
|            | Stage 3: Analysis of Protein Expression in Transfected Cells   | 17.68          |
|            | <ul> <li>Alternative Protocol: Tetracycline-regulated Induction of Gene Expression in<br/>Transiently Transfected Cells Using the Autoregulatory tTA System</li> </ul> | 17.70          |
|            | 9 Ecdysone as Regulator of Inducible Gene Expression in Mammalian Cells  | 17.71          |
| 11         | NFORMATION PANELS  |                |
|            | Footprinting DNA   | 17.75          |
|            | Gel Retardation Assays   | 17.78          |
|            | Baculoviruses and Baculovirus Expression Systems   | 17.81          |
|            | Green Fluorescent Proteins   | 17.84          |
|            | Epitope Tagging  | 17.90          |
|            | Chloramphenicol Acetyltransferase  | 17.94          |
|            | Luciferase   | 17.96          |
|            | β-galactosidase  | 17.97          |
| Chapter 18 |  |                |
| •          | tion Technologies  | 18.1           |
| 110        | NTRODUCTION  |                |
| P          | ROTOCOLS   |                |
| Ţ.         | 1 Two-hybrid and Other Two-component Systems   | 18.6           |
|            | Stage 1: Characterization of a Bait-LexA Fusion Protein  | 18.17          |
| V          | Alternative Protocol: Assay of β-galactosidase Activity by Chloroform Overlay  | 18.28          |
|            | Stage 2: Selecting an Interactor   | 18.30          |
|            | Stage 3: Second Confirmation of Positive Interactions  | 18.38          |
|            | Alternative Protocol: Rapid Screen for Interaction Trap Positives  | 18.46          |
|            | 2 Detection of Protein-Protein Interactions Using Far Western with GST Fusion Proteins   | 18.48          |
|            | <ul> <li>Additional Protocol: Refolding of Membrane-bound Proteins</li> </ul>  | 18.53          |
|            | <ul> <li>Alternative Protocol: Detection of Protein-Protein Interactions with Anti-GST<br/>Antibodies</li> </ul>   | 18.54          |
|            | 3 Detection of Protein-Protein Interactions Using the GST Fusion Protein Pulldown<br>Technique   | 18.55          |
|            | 4 Identification of Associated Proteins by Coimmunoprecipitation   | 18.60          |
|            | 5 Probing Protein Interactions Using GFP and Fluorescence Resonance Energy<br>Transfer   | 18.69          |
|            | Stage 1: Labeling Proteins with Fluorescent Dyes   | 18.80          |

#### Contents

| 18.8        |
|-------------|
| ysis 18.8   |
| 18.8        |
| 18.9        |
| oscopy 18.9 |
|             |
| 18.10       |
| 18.10       |
|             |

#### INFORMATION PANELS

| Filamentous Phage Display                 | 18.115 |
|---|--------|
| Genomics and the Interaction Trap         | 18.123 |
| Interaction Trap and Related Technologies | 18.125 |

### Appendices

- 1 Preparation of Reagents and Buffers Used in Molecular Cloning, A1.1
- 2 Media, A2.1
- 3 Vectors and Bacterial Strains, A3.1
- 4 Enzymes Used in Molecular Cloning, A4.1
- 5 Inhibitors of Enzymes, A5.1
- 6 Nucleic Acids, A6.1
- 7 Codons and Amino Acids, A7.1
- 8 Commonly Used Techniques in Molecular Cloning, A8.1
- 9 Detection Systems, A9.1
- 10 DNA Array Technology, A10.1
- 11 Bioinformatics, A11.1
- 12 Cautions, A12.1
- 13 Suppliers, A13.1
- 14 Trademarks, A14.1
  - Appendix References, R1

INDEX, 1,1